

## Induction of Labour

### Making your choice

Everyone has the right to be fully informed and to share in decision-making about their health care. Before you make a decision about induction, your doctor or midwife will explain:

- why an induction has been recommended for you, and the potential benefits
- the potential risks with continuing your pregnancy until labour starts naturally
- potential risks with having an induction of labour
- the procedures and care that is involved during an induction of labour.

Some women will choose to 'wait and see' whether natural labour will start. However it is important that you are aware of the risks of both options, so that you can decide what is best for you.

### What is induction of labour?

In most pregnancies, labour starts naturally between 37 and 42 weeks, leading to the birth of the baby. When labour starts, a number of changes take place in your body:

- the cervix (neck of the womb) softens and shortens
- the fluid-filled membrane sac surrounding your baby tears ('your waters break')
- the cervix dilates (opens)
- the womb contracts to push your baby out.

### When is induction of labour recommended?

Approximately one fifth of women have an induction of labour. The most common reasons are:

- the woman has specific health concerns (such as diabetes or high blood pressure)
- the baby is not well or is distressed
- the pregnancy has gone longer than 41 weeks (prolonged pregnancy)
- the waters have already broken but the contractions of labour have not started naturally.

An induction is recommended when it is considered that your health and/or your baby's health will benefit.

### How is labour induced?

Before starting the induction, your doctor or midwife will assess your cervix (the neck of the womb). This examination takes only a few minutes but some women may experience some discomfort. Based on this examination your doctor or midwife will recommend one of the following methods of induction:

- prostaglandin
- cervical ripening balloon catheter
- oxytocin
- Artificially breaking the waters.

Induction can be one or a combination of these methods.

#### Risks / things you should be aware of:

- Induction for reasons other than prolonged pregnancy
- May increase the chance of you having a caesarean section.
- Women who are induced are more likely to experience above average blood loss after the birth.
- In the event the birth suites are busy, your induction of labour may be delayed and the process of induction may take longer than one day.

**All women and their partners are encouraged to ask questions and express any concerns or wishes they may have at any stage during the planning and process of induction of labour.**

### Prostaglandin

Prostaglandin is a naturally occurring hormone that prepares your body for labour. A synthetic version has been developed to mimic the effect of the hormone. This is inserted into your vagina, usually in the form of a gel which slowly releases the prostaglandin over 12–24 hours. When the prostaglandin is in place, you will have a CTG monitoring for least 1 hour after the gel is inserted. Once the prostaglandin has been inserted you will need to remain in hospital. You should inform your midwife immediately if you experience any of the following:

- regular painful contractions 5 mins apart for your first baby, or 10 mins apart for subsequent babies
- your membranes rupture (your waters break) spontaneously

- your baby seems to be moving less
- you have vaginal bleeding.

When the prostaglandin takes effect, your cervix will soften and open. If the gel is used, you may require one, two, or three doses (given every six to eight hours). When the cervix is soft and open, your body is prepared for labour. The next steps will vary from woman to woman – some might require an ARM to ‘break their waters’, whereas this might happen naturally for other women. Some women might require oxytocin to stimulate the contractions.

#### **Risks / things you should be aware of**

- Prostaglandin sometimes causes vaginal soreness. However, there is no evidence to suggest that labour induced with prostaglandin is any more painful than labour that has started naturally.

#### **Cervical ripening balloon catheter**

Prostaglandin does not suit all women and there will be circumstances in which your doctor may recommend using a cervical ripening balloon catheter. This catheter is inserted into your cervix and the balloons inflated with saline, thus applying pressure to the cervix. The pressure should soften and open your cervix, thereby preparing your body for labour.

When the catheter is in place, you will need to stay in hospital but you will be able to move around normally. During this time the midwives will periodically check you and listen to your baby’s heart.

Please tell the midwife caring for you if:

- your catheter falls out
- you have regular painful contractions; 5 mins apart if this is your first baby, or 10 mins apart for subsequent babies
- your membranes rupture (your waters break)spontaneously
- your baby seems to be moving less
- You have vaginal bleeding.

What happens next will vary from woman to woman the catheter usually remain in for 12 – 14 hours – some might require an ARM to “break their waters”, whereas this might happen naturally for other women. Some women might require oxytocin to stimulate the contractions.

#### **Risks / things you should be aware of:**

- The vaginal examination needed to performance this procedure may cause you some discomfort.

#### **Artificial Rupture of Membranes (‘breaking your waters’)**

If your waters have not broken, a procedure called an ‘Artificial Rupture of Membranes’ or ‘ARM’ may be recommended. This is when your midwife or doctor makes a hole in your membrane sac to release the fluid inside. This procedure is done through your vagina using a small instrument. Sometimes releasing the waters is enough to ‘get things going’ and labour will commence. However, most women will also require the oxytocin drug as well (described above) to start the contractions.

#### **Risks / things you should be aware of:**

- The vaginal examination needed to perform this procedure may cause you some discomfort.
- Although ARM is usually straightforward, it can increase the risk of cord prolapse, bleeding and infection.

#### **Oxytocin**

Oxytocin is the hormone that causes contractions. A synthetic version of oxytocin is given to women when contractions don’t start naturally. Oxytocin is given through a drip, and enters a vein in the arm. Once contractions begin, the rate of the drip is adjusted so that contractions occur regularly until your baby is born. This process can take several hours. Your baby’s heart rate will be monitored throughout labour using a CTG machine.

#### **Risks / things you should be aware of:**

- Your ability to move around will be limited by the drip and the CTG monitor. Whilst it may be okay to stand up or sit down, it will not be possible to have a bath or move from room to room.
- Very occasionally oxytocin can cause the uterus to contract too frequently which may affect the pattern of your baby’s heartbeat. If this happens you would be asked to lie on your left side and the drip will be slowed to lessen the contractions. Another drug may be given to counteract the oxytocin.

#### **Where to get more information**

Please phone the Maternity Unit prior to admission for their induction of labour as sometimes your induction need to be rescheduled.

**For further information please call the Maternity Unit on 56230 752**